

---

## Maize (*Zea mays* L.) Agro-Physiological Response to Potassium and Iron Fertilizer under Water Deficit Stress

Saeed Reza Yaghoobi<sup>1</sup>, Arash Roozbehani<sup>2\*</sup>, Mohammad Reza Akhavan Mohseni<sup>2</sup>

1- Assistant Professor, Technical University, Samangan Agricultural College, North Khorasan.

2- Department of Agronomy and Plant Breeding, Roudehen Branch, Islamic Azad University, Roudehen, Iran.

---

### RESEARCH ARTICLE

#### ARTICLE INFO.

*Received Date:* 3 Jul. 2017

*Received in revised form:* 4 Aug. 2017

*Accepted Date:* 2 Sep. 2017

*Available online:* 30 Sep. 2017

© 2015 IAUAHZ Publisher All Rights Reserved.

**To Cite This Article:** Saeed Reza Yaghoobi, Arash Roozbehani, Mohammad Reza Akhavan Mohseni. Maize (*Zea mays* L.) Agro-Physiological Response to Potassium and Iron Fertilizer under Water Deficit Stress. *J. Crop. Nut. Sci.*, 3(3): 37-48, 2017.

---

### ABSTRACT

This research was conducted to evaluate effect of potassium and iron fertilizers on agro-physiological traits affected different irrigation regime of corn in Shahryar (Tehran province, central of Iran) via a split-split plots arrangement based on randomized complete blocks design with three replications during 2015. The main plot included different irrigation regime (Normal irrigation and stop irrigation at grain filling period). Sub plot consisted different level of K<sub>2</sub>O (0, 25, 50 kg.ha<sup>-1</sup>), from water soluble potassium sulfate source. Sub-sub plot included Fe-EDDHA that apply as fertigation 2 kg.ha<sup>-1</sup>, foliar application Fe-EDTA 2 kg.ha<sup>-1</sup>, and no iron fertilizer application as control. Application of Fe-EDDHA as fertigation and Fe-EDTA as foliar application in normal irrigation increased corn seed yield from 5232 kg.ha<sup>-1</sup> in control to 6622 and 6464 kg.ha<sup>-1</sup> respectively but in water deficit situation were not effective. In normal irrigation, application 25 and 50 kg.ha<sup>-1</sup> soluble potassium sulfate increased corn seed yield from 5294 kg.ha<sup>-1</sup> in control to 6975 and 6048 kg.ha<sup>-1</sup> respectively. Under water deficit stress application of 25 and 50 kg.ha<sup>-1</sup> soluble potassium sulfate increased corn seed yield from 3921 kg.ha<sup>-1</sup> to 4794 and 4807 kg.ha<sup>-1</sup> respectively. Maximum corn seed yield was achieved when 25 kg.ha<sup>-1</sup> soluble potassium sulfate and Fe-EDDHA applied together in 6392 kg.ha<sup>-1</sup>. Application soluble potassium sulfate and iron fertilizers increased chlorophyll a, b, and total content in different irrigation regimes. However fertigation of soluble potassium sulfate along with Fe-EDDHA was more efficient to improve chlorophyll a, b, and total and seed yield in water deficit situation.

---

**Key words:** *Corn, Fertigation, Foliar application, Seed yield*

---

## INTRODUCTION

Water is about to become increasingly limited for crop production (Martineau *et al.*, 2017). Maize is the one of the important field crops in creating human food security (Afarinesh, 2015; Shiri *et al.*, 2015). Drought is one of the factors which threaten agriculture products in most parts of the world (Abolhasani and Saeidi, 2004; Banziger *et al.*, 2002). It causes stress in plants and is not only caused by the reduction of rainfalls and great heat, but in the cases where there is moisture in the soil, this moisture cannot be used for plants for some reasons such as excessive soil salinity or soil frost, and plants will be stressed (Baydar and Erbas, 2005). Drought and water shortage are considered an objective reality. In the past, water crisis was not as significant as today, since the population was less, but with the population increase by about six times and the need for more food during the last 100 years, the incidence of this crisis has become more evident than the past (Chimenti *et al.*, 2002). Production of maize for forage or seed requires a lot of water. However in last decades breeders focused on introducing drought tolerant crops (Tabatabaei and Shaleri, 2015) but plant nutrition got a special consideration in water deficit situation (Norastehnia and Farjadi, 2016). Between plant nutrition elements, potassium has a special value in biotic and abiotic stresses like drought stress. Potassium has many physiological role in plants such as enzymes activation in biological compounds synthesis, plant water relationship and stomates opening, photosynthesis and biomaterial transport, induction plant response to biotic and abiotic stresses (Oosterhuit *et al.*, 2013). There have been reports of positive effects of potassium on drought stress (Martineau *et al.*, 2017), salinity stress (Ullah Jan *et*

*al.*, 2017), chilling stress (Karimi, 2017), fungal disease (Zimmerman Lax *et al.*, 2018). Research has shown potassium availability in plants under drought stress will increase water absorption by root cells and root extending in soil improve its access to new water and nutrition contents (Grzebisz *et al.*, 2013). This reduces effects of drought stress on plant (Zahoor *et al.*, 2017) and prevents senescence (Lv *et al.*, 2017). Potassium is not only an essential macronutrient for plant growth and development, but also is a primary osmotic in maintaining low water potential of plant tissues. Therefore, for plants growing in drought conditions, accumulating abundant  $K^+$  in their tissues may play an important role in water uptake along a soil-plant gradient. In general,  $K^+$  is accumulated in response to soil water deficits, while  $Na^+$  is accumulated under saline conditions (Glenn *et al.*, 1996). Amalia-Muhd-Zain and Razi-Ismail (2016) reported application potassium in potassium chloride and potassium sulfate forms eliminated water deficit stress on rice growth and development. They emphasized potassium chloride was more effective than potassium sulfate in water deficit elimination. There are some researches that confirm potassium efficacy in subtraction of drought stress in field crop such as safflower (Abedi Baba- Arabi *et al.*, 2012), canola (Fanaei *et al.*, 2011), sesame (Aien, 2012), wheat (Ramezanzpour *et al.*, 2009), tobacco (Norastehnia and Farjadi, 2016) and sorghum (Khezerloo *et al.*, 2010). Iron is a micro nutrient element which has especial role in plant photosynthesis. But because of high content of carbonate and bicarbonate in soils from Iran, iron deficiency is obvious in sensitive crops (Martinez-Cuenca *et al.*, 2013). Furthermore water deficit stress complicated the problem. How-

ever application of chelated iron fertilizer somewhat solve the problem. Movahhedi Dehnavi and Jalil Sheshbahre (2017) showed that photosystem II of soybean leaves impressed by drought stress and photosynthesis reduced but application of iron and zinc fertilizers retrieved it. Also Ahmadi *et al.* (2016) obtained similar results about iron fertilizer on chick pea photosynthesis under drought stress. Goleg *et al.* (2016) indicated foliar application of ferrous nano oxide one  $\text{kg}\cdot\text{ha}^{-1}$  prevented the effects of drought stress on sesame. Dindoost and Yousefzadeh (2014) found that foliar application of iron fertilizer increased sunflower yield and oil percent under drought stress. Climate change caused long term droughts in some regions of world and agricultural products have been affected extensively. Under water shortage conditions, the effectiveness of fertilizers decreases, especially if consumption of these fertilizers is not compatible with the vegetative growth of plants. However, it should be noted that soils of Iran, which are categorized under the calcareous soils, due to drought stress, salinity, being calcareous, highly acidity, having low contents of organic materials, continuing drought, and continuing unbalanced consumption of fertilizers, iron and potassium contents are too low. Therefore, the plants which grow in such soils are mainly suffered from shortage of iron and potassium and shortage indications are observed in them (Jaleel *et al.*, 2009). However water supply for agriculture is getting harder every year but it is possible to produce satisfactory yield with a proper nutrient management under limit water resources. Main goal of research is investigation of potassium and iron fertilizers application on corn physiological properties under water deficit stress.

## MATERIALS AND METHODS

### *Field and Treatment Information*

This research was conducted to evaluate effect of potassium and iron fertilizers on agro-physiological traits affected different irrigation regime in Shahryar (Tehran province, central of Iran) via a split split plots arrangement based on randomized complete blocks design with three replications during 2015. Soil texture of research field was sandy loam with pH 7.8. Soil analysis results are shown in table 1. The main plot included different irrigation regime (Normal irrigation and stop irrigation at grain filling period). Sub plot included different level of  $\text{K}_2\text{O}$  (0, 25, 50  $\text{kg}\cdot\text{ha}^{-1}$ ), from water soluble potassium sulfat (Solupotasse, Tessenderlo Group, Belgium) source. Sub-sub plot included Fe-EDDHA [Fe- Ethylene di amine-N, N'-bis (2-hydroxy phenyl acetic acid)] that apply as fertigation 2  $\text{kg}\cdot\text{ha}^{-1}$ , foliar application Fe-EDTA [Fe Ethylenediaminetetra acetic acid] 2  $\text{kg}\cdot\text{ha}^{-1}$ , and no iron fertilizer application as control.

**Table 1.** Result of soil chemical analysis

TNV (%)	12
N (%)	2.66
$\text{P}_2\text{O}_5$ (ppm)	0.22
$\text{K}_2\text{O}$ (ppm)	10
MgO (ppm)	2.8
Fe (ppm)	0.16
Zn (ppm)	0.9

### *Farm Management*

Soil preparation was done with plowing and disking operation. Corn seeds SC704 variety was sown with 75 cm inter row distances with density of 10000 plants per hectare. Experimental plots included four cultured row with 5 m length. The irrigation was carried out at intervals five days and weeds were removed manually.

Potassium sulfate and Fe-EDDHA applied as fertigation and Fe-EDTA applied as foliar in early corn tasseling stage. Water deficit stress treatment was carried out by stop irrigation in milky seed filling stage of corn.

### Measured Traits

Chlorophyll a, b, total and carotenoid were measured in method by Arnon (1967). Leaf samples of corn obtained from nearest leaves to ear in milky seed filling stage. For pigments extract 0.5 g fresh corn leaf crushed in 10 ml acetone 80% solvent then its volume adjusted to 20 ml. The spectrophotometer was calibrated at zero absorption using a blank of pure solvent. Absorbance of samples measured at 645 and 663 nm no longer than 20 minutes after extraction. The equations 1, 2, 3 and 4 were used for chlorophyll and carotenoid calculation:

**Equ. 1.**  $\text{Chl.a (mg.g}^{-1}\text{ FWL)} = [12.7 (A_{663}) - 2.69 (A_{645})] V/1000 \times W$

**Equ. 2.**  $\text{Chl.b (mg.g}^{-1}\text{ FWL)} = [22.9 (A_{645}) - 4.68 (A_{663})] V/1000 \times W$

**Equ. 3.**  $\text{Chl.total (mg.g}^{-1}\text{ FWL)} = \text{Chl.a} + \text{Chl.b}$

**Equ. 4.**  $\text{Carotenoid} = [1000 (A_{470}) - 1.8 (\text{Chl.a}) - 85.02 (\text{Chl.b})] \times 20$

Chl.a: Chlorophyll a, Chl.b: Chlorophyll b, FWL: Fresh weight leaf, A = absorbance at specific wave length, V= final volume of chlorophyll extract in 80 % acetone. W= fresh weight of tissue extract.

At corn harvest, considering the marginal effects, two meter squares of each plot harvested and seed and biological yield measured. Harvest index calculated with equation 5 (Donald and Hamblin, 1976):

**Equ. 5.**  $\text{Harvest index} = (\text{Seed yield} / \text{Biological yield}) \times 100$

### Statistical Analysis

Statistical analysis (ANOVA) was applied with using the by the SAS software (Ver.9.1). The differences between

treatment means were assessed by least significant difference (LSD) test at 5% probability level.

## RESULT

### Chlorophyll a

Result of analysis of variance showed that the effect of different levels of irrigation regime, potassium and iron fertilizer and interaction effect of treatments (instead irrigation regime  $\times$  potassium  $\times$  iron) on chlorophyll a was significant at 1% probability level (Table 2). Mean comparison result of interaction effect of different level of irrigation regime and potassium fertilizer indicated that maximum chlorophyll a ( $2.70 \text{ mg.g}^{-1}$ ) was noted for normal irrigation with use  $25 \text{ kg.ha}^{-1} \text{ K}_2\text{O}$  and minimum of that ( $1.40 \text{ mg.g}^{-1}$ ) belonged to water stress with non-use potassium fertilizer treatment (Fig.1a). As for LSD classification made with respect to interaction effect of different level of irrigation regime and iron fertilizer maximum and minimum amount of chlorophyll a belonged to normal irrigation with use Fe-EDDHA ( $2.25 \text{ mg.g}^{-1}$ ) and water stress with non-use iron fertilizer ( $1.60 \text{ mg.g}^{-1}$ ) (Fig.1b). According mean comparison result of interaction effect of different level of potassium and iron fertilizer maximum amount of chlorophyll a ( $2.30 \text{ mg.g}^{-1}$ ) was obtained for  $25 \text{ kg.ha}^{-1} \text{ K}_2\text{O}$  with use Fe-EDDHA and minimum of that ( $1.38 \text{ mg.g}^{-1}$ ) was for control treatment (Fig.1c).

### Chlorophyll b

According result of analysis of variance the effect of different levels of irrigation regime, potassium and iron fertilizer and interaction effect of treatments (instead irrigation regime  $\times$  potassium  $\times$  iron) on chlorophyll b was significant at 1% probability level (Table 2).

Evaluation mean comparison interaction effect of different level of irrigation regime and potassium fertilizer revealed the maximum amount of chlorophyll b ( $1.26 \text{ mg.g}^{-1}$ ) was noted for normal irrigation with use  $25 \text{ kg.ha}^{-1} \text{ K}_2\text{O}$  and minimum of that ( $0.48 \text{ mg.g}^{-1}$ ) belonged to water stress with non-use potassium fertilizer treatment (Fig.2a). Assessment result of interaction effect of different level of irrigation regime and iron fertilizer showed the maximum chlorophyll b ( $1.14 \text{ mg.g}^{-1}$ ) was noted for normal irri-

gation with use Fe-EDDHA and minimum of that ( $0.47 \text{ mg.g}^{-1}$ ) belonged to water stress with non-use iron fertilizer (Fig.2b). Evaluation mean comparison result of interaction effect of different level of potassium and iron fertilizer indicated maximum amount of chlorophyll b ( $1.17 \text{ mg.g}^{-1}$ ) was noted for  $25 \text{ kg.ha}^{-1} \text{ K}_2\text{O}$  with use Fe-EDDHA and lowest one ( $0.44 \text{ mg.g}^{-1}$ ) belonged to control treatment (Fig.2c).

**Table 2.** Result of Analysis of variance of the effect of different level of irrigation regime, potassium and iron fertilizer on measured traits


S.O.V	df	Chlorophyll a	Chlorophyll b	Total chlorophyll	Carotenoid to chlorophyll ratio	Seed yield
<b>Block</b>	2	0.0097 <sup>ns</sup>	0.018 <sup>ns</sup>	0.051 <sup>ns</sup>	0.00027 <sup>ns</sup>	107725 <sup>ns</sup>
<b>Irrigation (I)</b>	1	3.27 <sup>**</sup>	2.17 <sup>**</sup>	10.77 <sup>**</sup>	0.24 <sup>*</sup>	25100698 <sup>**</sup>
<b>Main Error</b>	2	0.018	0.024	0.088	0.0004	184545
<b>Potassium (P)</b>	2	0.59 <sup>**</sup>	0.51 <sup>**</sup>	2.18 <sup>**</sup>	0.0014 <sup>ns</sup>	3846645 <sup>**</sup>
<b>I×P</b>	2	0.41 <sup>**</sup>	0.43 <sup>**</sup>	1.72 <sup>**</sup>	0.00011	2622892 <sup>**</sup>
<b>Sub Plot Error</b>	8	0.026	0.027	0.1	0.00034	108471
<b>Iron (Ir)</b>	2	0.51 <sup>**</sup>	0.37 <sup>**</sup>	1.77 <sup>**</sup>	0.0029 <sup>ns</sup>	3679020 <sup>**</sup>
<b>I × Ir</b>	2	0.29 <sup>**</sup>	0.26 <sup>**</sup>	1.12 <sup>**</sup>	0.00067 <sup>ns</sup>	1770283 <sup>**</sup>
<b>P × Ir</b>	4	0.16 <sup>**</sup>	0.17 <sup>**</sup>	0.65 <sup>**</sup>	0.00012 <sup>ns</sup>	1045386 <sup>**</sup>
<b>I × P × Ir</b>	4	0.165 <sup>ns</sup>	0.158 <sup>ns</sup>	0.67 <sup>ns</sup>	0.0012 <sup>ns</sup>	919120 <sup>ns</sup>
<b>Sub Sub Plot Error</b>	24	0.047	0.04	0.17	0.00089	290065
<b>CV (%)</b>	-	11.9	28	16.4	25	9.9

<sup>ns</sup>, \* and \*\* are non-significant and significant at 5 and 1% probability levels, respectively.

### Total chlorophyll

According result of analysis of variance the effect of different levels of irrigation regime, potassium and iron fertilizer and interaction effect of treatments (instead irrigation regime × potassium × iron) on total chlorophyll was significant at 1% probability level (Table 2). Assessment mean comparison interaction effect of different level of irrigation regime and potassium fertilizer indicated the maximum rate of total chlorophyll ( $3.69 \text{ mg.g}^{-1}$ ) was noted for normal irrigation with use  $25 \text{ kg.ha}^{-1} \text{ K}_2\text{O}$  and lowest one ( $2.02 \text{ mg.g}^{-1}$ ) belonged to water stress with non-use potassium fertilizer treatment (Fig.3a).

Evaluation result of interaction effect of different level of irrigation regime and iron fertilizer revealed the maximum rate of total chlorophyll ( $3.46 \text{ mg.g}^{-1}$ ) was noted for normal irrigation with use Fe-EDDHA and minimum of that ( $1.99 \text{ mg.g}^{-1}$ ) belonged to water stress with non-use iron fertilizer (Fig.3b). According mean comparison result of interaction effect of different level of potassium and iron fertilizer the maximum rate of total chlorophyll ( $3.46 \text{ mg.g}^{-1}$ ) was noted for  $25 \text{ kg.ha}^{-1} \text{ K}_2\text{O}$  with use Fe-EDDHA and the lowest one ( $1.90 \text{ mg.g}^{-1}$ ) belonged to the control treatment (Fig.3c).



**Fig. 1.** Mean comparison interaction effect of potassium fertilizer and irrigation regime (A), iron fertilizers and irrigation regime (B), potassium and iron fertilizers iron fertilizer (C) on chlorophyll a concentration via LSD test at 5% probability level.


### Carotenoid to chlorophyll ratio

Result of ANOVA showed only effect of different levels of irrigation regime on carotenoid to chlorophyll ratio was significant at 5% probability level (Table 2). Compare different irrigation regime revealed maximum carotenoid to chlorophyll ratio (0.14) was observed in water stress and the lowest one (0.1) was found in normal irrigation (Fig.4).


### Seed yield

According result of analysis of variance the effect of different levels of ir-

rigation regime, potassium and iron fertilizer and interaction effect of treatments (instead irrigation regime  $\times$  potassium  $\times$  iron) on seed yield was significant at 1% probability level (Table 2). Mean comparison result of interaction effect of different level of irrigation regime and potassium fertilizer indicated that maximum seed yield (6975 kg.ha<sup>-1</sup>) was noted for normal irrigation with use 25 kg.ha<sup>-1</sup> K<sub>2</sub>O and minimum of that (3921 kg.ha<sup>-1</sup>) belonged to water stress with non-use potassium fertilizer treatment (Fig.5a).




**Fig. 2.** Mean comparison interaction effect of potassium fertilizer and irrigation regime (A), iron fertilizers and irrigation regime (B), potassium and iron fertilizers iron fertilizer (C) on chlorophyll b concentration via LSD test at 5% probability level.



**Fig. 3.** Mean comparison interaction effect of potassium fertilizer and irrigation regime (A), iron fertilizers and irrigation regime (B), potassium and iron fertilizers iron fertilizer (C) on total chlorophyll concentration via LSD test at 5% probability level.


With respect to interaction effect of different level of irrigation regime and iron fertilizer maximum and minimum seed yield belonged to normal irrigation with use Fe-EDDHA (6622 kg.ha<sup>-1</sup>) and water stress with non-use iron fertilizer (4572 kg.ha<sup>-1</sup>) (Fig.5b). According mean comparison result of interaction effect of potassium and iron fertilizer maximum amount of seed yield (6392 kg.ha<sup>-1</sup>) was obtained for 25 kg.ha<sup>-1</sup> K<sub>2</sub>O with use Fe-EDDHA and minimum of that (4402 kg.ha<sup>-1</sup>) was for control treatment (Fig.5c).



**Fig. 4.** Mean comparison different level of irrigation regime on carotenoid to chlorophyll via LSD test at 5% probability level.

## DISCUSSION

Application of soluble potassium sulfate in normal irrigation significantly increased chlorophyll a, b and total concentration. Total chlorophyll concentration of corn reduced significantly when irrigation stopped in tasseling from 2.36 to 1.99 mg.g<sup>-1</sup> leaf fresh weight (Fig. 1a, 2a and 3a) like chlorophyll a and b. Application of Fe-EDDHA and Fe-EDTA caused maximum chlorophyll a, b and total concentrations in normal irrigation condition but in water deficit condition there was no significant difference between iron fertilizer application and control treatment (Fig. 1b, 2b and 3b). Maximum total chlorophyll concentration was 3.69 mg.g<sup>-1</sup> leaf fresh weights when soluble potassium sulfate used in 25 kg.ha<sup>-1</sup>. In water deficit condition application soluble potassium sulfate changed chlorophyll concentration 2.02 to 2.12 mg.g<sup>-1</sup> corn leaf fresh weight which there was no significance (Fig. 3a). It was happened for chlorophyll a and b too (Fig. 1a, 2a). As results application of iron fertilizer increased chlorophyll concentration from 1.9 mg.g<sup>-1</sup> corn leaf fresh weight to 2.36 mg.g<sup>-1</sup> corn leaf fresh weight in Fe-EDDHA fertigation and 2.37 mg.g<sup>-1</sup> corn leaf fresh weight in Fe-EDTA foliar application under no soluble potassium sulfate application.



**Fig. 5.** Mean comparison interaction effect of potassium fertilizer and irrigation regime (A), iron fertilizers and irrigation regime (B), potassium and iron fertilizers iron fertilizer (C) on seed yield concentration via LSD test at 5% probability level.

Application 25 kg.ha<sup>-1</sup> increased total chlorophyll concentration from 2.17 mg.g<sup>-1</sup> corn leaf fresh weight to 3.46 mg.g<sup>-1</sup> corn leaf fresh weight in Fe-EDDHA fertigation and 3.08 mg.g<sup>-1</sup> corn leaf fresh weight in Fe-EDTA foliar application (Fig. 3c). The similar results obtained for chlorophyll a and b (Fig. 1c, 2c). However potassium and iron fertilizers application increased chlorophyll a, b and total in corn leaves in normal irrigation but in water deficit stress which was interrupted irrigation in corn milky seed filling stage did not

changed chlorophyll concentration. Lv *et al.* (2017) showed potassium application after pollination stage increased chlorophyll concentration in wheat. Asgharipour and Heidari (2011) reported sulfate potassium application increased sorghum leaves chlorophyll content. They revealed potassium availability in soil caused drought resistance in sorghum. As shown in Fig. 4, in normal irrigation ratio of carotenoid to chlorophyll was 0.1 but under water deficit situation the ratio of carotenoid to chlorophyll obtained 0.14. It seems that the chlorophyll content of corn leaves is more affected by water deficit stress than carotenoids (Fig. 4). In normal irrigation, application 25 and 50 kg.ha<sup>-1</sup> soluble potassium sulfate increased corn seed yield from 5294 kg.ha<sup>-1</sup> in control to 6975 and 6048 kg.ha<sup>-1</sup> respectively. Under water deficit stress application of 25 and 50 kg.ha<sup>-1</sup> soluble potassium sulfate increased corn seed yield from 3921 kg.ha<sup>-1</sup> to 4794 and 4807 kg.ha<sup>-1</sup> respectively (Fig. 5a). Application of Fe-EDDHA as fertigation and Fe-EDTA as foliar application in normal irrigation increased corn seed yield from 5232 kg.ha<sup>-1</sup> in control to 6622 and 6464 kg.ha<sup>-1</sup> respectively but in water deficit situation were not effective (Fig. 5b). Maximum corn seed yield was achieved when 25 kg.ha<sup>-1</sup> soluble potassium sulfate and Fe-EDDHA applied together in 6392 kg.ha<sup>-1</sup> (Fig. 5c). In present world water deficiency is one the important problems especially in agriculture. Due to direct effect on root, water deficit stress has influence adversely on nutrient absorption by plant (Wang *et al.*, 2013). However proper nutrition program could reduce negative effects. Premachandra *et al.* (1991) reported availability of potassium increases corn resistance to drought stress because of osmotic balance improvement.



In present study potassium application along with normal irrigation have positive effects on chlorophyll a, b and total content and seed yield of corn. In water deficit situation however potassium application had positive effects on mentioned traits but were not significant. Potassium has important role in transport and metabolism of poly saccharides and amino acids on sink (Hu *et al.*, 2018) and sink content is directly concern to cereal seed filling (Lv *et al.*, 2017; Arif *et al.*, 2017). In other hand potassium is important in opening of stomata and adaptation to surrounding environment (Hasanuzzaman *et al.*, 2018) and leaf long life (Lv *et al.*, 2017). There are similar reports about potassium important role in cotton (Zahoor *et al.*, 2017), rice (Chen *et al.*, 2017) corn (Aslam *et al.*, 2013), and sunflower (Soleimanzadeh *et al.*, 2010). Under normal irrigation in present study application iron fertilizer to both Fe-EDDHA as fertigation and Fe-EDTA as foliar, had positive effects on chlorophyll a, b and total and seed yield of corn. Caliskan *et al.* (2008) revealed application of 400 gr.ha<sup>-1</sup> Fe-EDTA along with 80 kg.ha<sup>-1</sup> nitrogen had the best soybean yield. They emphasized application of coated nitrogen fertilizer along with Fe-EDTA as starter increased early plant growth. Janmohammadi *et al.* (2018) reported application of iron and zinc fertilizers along with manure had good results on pea yield. Like this maximum sweet corn seed yield obtained when enriched vermin compost with iron sulfate and zinc sulfate along with foliar application of zinc and iron sulfate applied 20 to 40 days after planting (Arabhanvi and Hulihalli, 2018). In present study however water deficit study reduced corn seed yield but iron fertilizer application somehow compensate the problem. Indeed Heidari *et al.* (2011) indicated application

of iron and sulfur fertilizers in water deficit stress affected significantly on iron and nitrogen content and yield of sesame.

## CONCLUSION

However water deficit stress is one of the major problems for crop production in present world but it could diminish somewhat by a proper plant nutrition program. As results of present study soluble potassium sulfate application along with Fe-EDDHA in milky seed filling could reduce somewhat effect of water deficit stress on corn.

## REFERENCES

- Abedi Baba-Arabi, S., M. Movahhedi Dehnavi, A. R. Yadavi. and E. Adhami. 2012.** Effects of Zn and K foliar application on physiological traits and yield of spring safflower under drought stress. *E. J. Crop Prod.* 4(1): 75-95. (Abstract in English)
- Abolhasani K. and G. Saeidi. 2004.** Relationships between agronomic characteristic of safflower under water stress and control. *Iranian J. Field Crop Res.* 1: 127-138. (Abstract in English)
- Afarinesh, A., G. R. Fathi, Chugan, S. A. Syadat, Kh. Alamisaid. and S. R. Ashrafizadeh. 2015.** Effects of drought stress and soil compaction on some agronomic traits of maize. *J. Plant Prod.* 38(2): 13-24. (Abstract in English)
- Ahmadi, L., M. Ghobadi, A. Saeidi. and J. Ghaderi. 2016.** The study of some physiologic and biochemical traits to drought stress and methods of Fe fertilizer application in chickpea. *J. Plant Process Function.* 4(14): 165-176. (Abstract in English)
- Aien, A. 2012.** Proline and soluble carbohydrate contents and potassium, zinc calcium uptake in sesame cultivars under drought stress. *J. Crop Prod. Environ. Stress.* 4(3): 39-48. (Abstract in English)

- Amalia-Mohd-Zain, N. and M. Razi Ismail. 2016.** Effects of potassium rates and types on growth, leaf gas exchange and biochemical changes in rice (*Oryza sativa*) planted under cyclic water stress. *Agric. Water Manage.* 164(1): 83-90.
- Arabhanvi, F. and U. K. Hulihalli. 2018.** Agronomic fortification with zinc and iron to enhancing micronutrient concentration in sweet corn seed to ameliorate the deficiency symptoms in human beings. *Intl. J. Current Micro-Biol. Appl. Sci.* 7(2): 333-340.
- Arnon, A. N. 1967.** Method of extraction of chlorophyll in the plants. *Agron. J.* 23: 112-121.
- Arif, M., M. Tasneem, F. Bashir, G. Yahseen. and A. Anwar. 2017.** Evaluation of different levels of potassium and zinc fertilizer on growth and yield of wheat. *Intl. J. Bio-Sensors Bio-Electronics.* 3(2): 57- 62.
- Asgharipour, M. R. and M. Heidari. 2011.** Effect of potassium supply on drought resistance in sorghum: plant growth and macronutrient content. *Pakistan J. Agric. Sci.* 48(3): 197-204.
- Aslam, M., M. S. I. Zamir, I. Afzal, M Yaseen, M. Mubeen. and A. Shoaib. 2013.** Drought stress, its effect on maize production and development of drought tolerance through potassium application. *Cercetari Agronomice in Moldova.* 2(154): 99-114.
- Banziger, M., G. O. Edmeades. and H. R. Lafitte. 2002.** Physiological mechanisms contributing to the increased N stress tolerance of tropical maize selected for drought tolerance. *Field Crops Res.* 75(3): 223-233.
- Baydar, H. and S. Erbas. 2005.** Influence of seed development and seed position on oil, fatty acids and total tocopherol contents in sunflower (*Helianthus annuus* L.). *Turkish J. Agric.* 29: 179-186.
- Chimenti, C. A., J. Pearson. and A. J. Hall. 2002.** Osmotic adjustment and yield maintenance under drought in sunflower. *Field Crop Res.* 75: 235-246.
- Caliskan, S., L. Ozkaya, M. E. Caliskan. and M. Arslan. 2008.** The effects of nitrogen and iron fertilization on growth, yield and fertilizer use efficiency of soybean in a Mediterranean-type soil. *Field Crop Res.* 108(2): 126-132.
- Chen, G., C. Liu, Z. Gao, Y. Zhang, H. Jiang, L. Zhu, D. Ren, L. Yu, G. Xu. and Q. Qian. 2017.** OsHAK1, a High-Affinity potassium transporter, positively regulates responses to drought stress in rice. *J. Frontiers in Plant Sci.* 8: 1-17.
- Dindoost, S. and S. Yousefzadeh. 2014.** Zinc, iron and manganese fertilizers foliar application on sunflower (*helianthus annuus* L., Var. Haisan) under drought stress. *J. Res. Agron. Sci.* 6(22): 25-41. (Abstract in English)
- Donald, C. M. and J. Hamblin. 1976.** The Biological yield and harvest index of cereals as agronomic and plant breeding criteria. *J. Adv. Agron.* 28: 361-405.
- Fanaei, H. R., M. Galavi, M. Kafi, A. Ghanbari Bonjar. and A. H. Shirani-Rad. 2011.** Effect of drought stress and potassium on solutes accumulation and chlorophyll of canola and Indian mustard. *J. Water Soil Sci.* 15(57): 141-156. (Abstract in English)
- Glenn, E., R. Pfister, J. J. Brown, T. L. Thompson. and J. Oleary. 1996.** Na<sup>+</sup> and K<sup>+</sup> accumulation and salt tolerance of *Atriplex canescens* genotypes. *Am. J. Bot.* 83: 997-1005.
- Goleg, M., H. Ghorbani. and M. Baradarn Firozabadi. 2016.** Effect of drought stress and foliar application of iron oxide nanoparticles on yield, ion content and photosynthetic pigments in sesame. *Iranian J. Field Crop Sci.* 46(4): 619-628. (Abstract in English)

- Grzebisz, W., A. Gransee, W. Szczepaniak, and J. Diatta. 2013.** The effect of potassium fertilization on water use efficiency in crop plants. *J. Plant Nutr. Soil Sci.* 176(3): 355-374.
- Hasanuzzaman, M., M. H. M. B. Bhuyan, K. Nahar, M. S. Hossain, J. A. Mahmud, M. S. Hossen, A. A. C. Masud, L. Moumita, and M. Fujita. 2018.** Potassium: A vital regulator of plant responses and tolerance to abiotic stress. *Agron. J.* 8(3): 31-38.
- Heidari, M., M. Galavi, and M. Hasani. 2011.** Effects of sulfur and iron fertilizers on yield, yield components and nutrient uptake in sesame under water stress. *Afr. J. Bio-Tech.* 10(44): 8816-8822.
- Hu, W., D. A. Loka, T. R. Fitzsimons, Z. Zhou, and D. M. Oosterhuis. 2018.** Potassium deficiency limits reproductive success by altering carbohydrate and protein balances in cotton. *Environ. Exp. Bot.* 145: 87-94.
- Jaleel, C. A., P. Manivanannan, A. M. Wahid, H. J. Froog, R. Al-Juburi, and R. Somasundaram. 2009.** Drought stress in plant: A review on morphological characters and pigments composition. *Intl. J. Agric. Biol.* 11: 100-105.
- Janmohammadi, M., H. Abdoli, N. Sabaghnia, M. Esmailpour, and A. Aghaei. 2018.** The effect of iron, zinc and organic fertilizer on yield of chickpea in Mediterranean climate. *Acta Universitatis Agriculturae Silviculturae Mendelianae Brunensis.* 66(1): 49-60.
- Karimi, R. 2017.** Potassium-induced freezing tolerance is associated with endogenous abscisic acid, polyamines and soluble sugars changes in grapevine. *Scientia Horticulturae.* 215(24): 184-194.
- Khezerloo, F., F. Jalili, and J. Khalili Mahaleh. 2010.** Drought stress and nitrogen and potassium on sorghum (Var. Speed feed) forage production. *J. Res. Agron. Sci.* 2(8): 51-66.
- Lv, X., T. Li, X. Wen, Y. Liao, and Y. Liu. 2017.** Effect of potassium foliage application post-anthesis on seed filling of wheat under drought stress. *Field Crops Res.* 206: 95-105.
- Martineau, E., J. C. Domec, A. Bosc, P. Denoroy, V. A. Fandino, J. Lavres, and L. Jordan-Meille. 2017.** The effects of potassium nutrition on water use in field-grown maize (*Zea mays* L.). *Environ. Exp. Bot.* 134: 62-71.
- Martinez-Cuenca, M. R., F. Legaz, M. A. Forner-Giner, E. Primo-Millo, and D. J. Iglesias. 2013.** Bicarbonate blocks iron translocation from cotyledons inducing iron stress responses in citrus roots. *J. Plant Physiol.* 170(10): 899-905.
- Movahhedi-Dehnavi, M. and M. Jalil Sheshbahre. 2017.** Soybean leaf physiological responses to drought stress improved via enhanced seed zinc and iron concentrations. *J. Plant Process and Function.* 5(18): 13-21. (Abstract in English)
- Norastehnia, A. and M. Farjadi. 2016.** The effect of the interaction between water stress and potassium nitrate on some of the physiological responses of *Nicotiana tabacum* L. *Nova Biologica Reperta.* 2(4): 260-271.
- Oosterhuit, D. M., D. A. Loka, and T. B. Raper. 2013.** Potassium and stress alleviation: Physiological functions and management of cotton. *Hournal Plant Nutr. Soil Sci.* 176(3): 331-343.
- Premachandra, G. S., H. Saneoka, and S. Ogata. 1991.** Cell membrane stability and leaf water relations as affected by potassium nutrition of water-stressed maize. *J. Exp. Bot.* 42(239): 739-745.
- Ramezanpour, M. R., M. Dastfal, and M. J. Malakouti. 2009.** Potassium effect on drought stress in wheat on Darab region, Fars province. *Iranian J. Soil Sci.* 22(1): 127-135. (Abstract in English)

- Shiri, M. R., R. Choukan. and R. T. Aliyev. 2015.** Drought stress effects on gene action and combining ability of maize inbred lines. *Seed Plant Improvement J.* 31(3): 421-440. (Abstract in English)
- Soleimanzadeh, H., D. Habibi, M. R. Ardakani, F. Paknejad. and F. Rejali. 2010.** Effect of potassium levels on antioxidant enzymes and malondialdehyde content under drought stress in sunflower (*Helianthus annuus* L.). *Am. J. Agric. Biol. Sci.* 5(1): 56-61.
- Tabatabaei, S. A. and E. Shaleri. 2015.** Short communication: effect of drought stress on maize hybrids yield and determination of the best hybrid using drought tolerance indices. *Environ. Stress Crop Sci.* 8(1): 121-125. (Abstract in English)
- Ullah Jan, A., F. Hadi, M. Midrarullah, M. Asif Nawaz. and K. Rahman. 2017.** Potassium and zinc increase tolerance to salt stress in wheat (*Triticum sativum* L.). *Plant Physiol. Bio-Chem.* 116: 139-149.
- Wang, M., Q. Zheng, Q. Shen. and S. Guo. 2013.** The critical role of potassium in plant stress response. *Intl. J. Molecular Sci.* 14(4): 7370-7390.
- Zahoor, R., W. Zhao, H. Dong, J. L. Snider, M. Abid, B. Iqbal. and Z. Zhou. 2017.** Potassium improves photosynthetic tolerance to and recovery from episodic drought stress in functional leaves of cotton. *Plant Physiol. Bio-Chem.* 119: 21-32.
- Zimmerman-Lax, N., D. Tamir- Ariel, M. Shenker. and S. Burdman. 2017.** Decreased potassium fertilization is associated with increased pathogen growth and disease severity caused by *Acidovorax citrulli* in melon foliage. *J. General Plant Pathol.* 84(1): 27-34.